

Surgical Treatments for Epilepsies in Children Aged 1–36 Months

A Systematic Review

Amy Y. Tsou, MD, MSc, Sudha Kilaru Kessler, MD, MSCE, Mingche Wu, MPH, Nicholas S. Abend, MD, MSCE, Shavonne L. Massey, MD, MSCE, and Jonathan R. Treadwell, PhD

Neurology® 2023;100:e1-e15. doi:10.1212/WNL.0000000000201012

Correspondence

Dr. Tsou
amytso@gmail.com

Abstract

Background and Objectives

Early life epilepsies (epilepsies in children 1–36 months old) are common and may be refractory to antiseizure medications. We summarize findings of a systematic review commissioned by the American Epilepsy Society to assess evidence and identify evidence gaps for surgical treatments for epilepsy in children aged 1–36 months without infantile spasms.

Methods

EMBASE, MEDLINE, PubMed, and the Cochrane Library were searched for studies published from 1/1/1999 to 8/19/21. We included studies reporting data on children aged 1 month to ≤36 months undergoing surgical interventions or neurostimulation for epilepsy and enrolling ≥10 patients per procedure. We excluded studies of infants with infantile spasms or status epilepticus. For effectiveness outcomes (seizure freedom, seizure frequency), studies were required to report follow-up at ≥ 12 weeks. For harm outcomes, no minimum follow-up was required. Outcomes for all epilepsy types, regardless of etiology, were reported together.

Results

Eighteen studies (in 19 articles) met the inclusion criteria. Sixteen prestudies/poststudies reported on efficacy, and 12 studies addressed harms. Surgeries were performed from 1979 to 2020. Seizure freedom for infants undergoing hemispherectomy/hemispherotomy ranged from 7% to 76% at 1 year after surgery. For nonhemispheric surgeries, seizure freedom ranged from 40% to 70%. For efficacy, we concluded low strength of evidence (SOE) suggests some infants achieve seizure freedom after epilepsy surgery. Over half of infants undergoing hemispherectomy/hemispherotomy achieved a favorable outcome (Engel I or II, International League Against Epilepsy I to IV, or >50% seizure reduction) at follow-up of >1 year, although studies had key limitations. Surgical mortality was rare for functional hemispherectomy/hemispherotomy and nonhemispheric resections. Low SOE suggests postoperative hydrocephalus is uncommon for infants undergoing nonhemispheric procedures for epilepsy.

Discussion

Although existing evidence remains sparse and low quality, some infants achieve seizure freedom after surgery and ≥50% achieve favorable outcomes. Future prospective studies in this age group are needed. In addition to seizure outcomes, studies should evaluate other important outcomes (developmental outcomes, quality of life [QOL], sleep, functional performance, and caregiver QOL).

Trial Registration Information

This systematic review was registered in PROSPERO (CRD42021220352) on March 5, 2021.

From the ECRI Evidence Based Practice Center (A.Y.T., M.W., J.R.T.), Plymouth Meeting, PA; Division of Neurology (A.Y.T.), Michael J Crescenz Veterans Affairs Medical Center, Philadelphia; Department of Pediatrics (Division of Neurology) (S.K.K., N.S.A., S.L.M.), Children's Hospital of Philadelphia; Departments of Neurology and Pediatrics (S.K.K., N.S.A., S.L.M.), University of Pennsylvania Perelman School of Medicine; Department of Anesthesia & Critical Care (N.S.A.), University of Pennsylvania Perelman School of Medicine; and Department of Biostatistics (N.S.A.), Epidemiology and Informatics, University of Pennsylvania Perelman School of Medicine.

Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article.

The Article Processing Charge was funded by the authors.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

RELATED ARTICLES

Editorial

Taming the Wild West of Early Life Epilepsy Care: One Question at a Time
Page 11

Research Article

Pharmacologic and Dietary Treatments for Epilepsies in Children Aged 1–36 Months: A Systematic Review
Page 17

Glossary

AES = American Epilepsy Society; **AHRQ** = Agency of Healthcare and Quality; **ASM** = anti-seizure medication; **DRE** = drug resistant epilepsy; **HME** = hemimegalencephaly; **ILAE** = International League Against Epilepsy; **MCD** = malformation of cortical development; **QOL** = quality of life; **SOE** = strength of evidence; **VPS** = ventriculoperitoneal shunt.

Onset of epilepsy in early life (age younger than 3 years) often has lifelong consequences. Seizures and disordered neuronal activity accompanying epilepsy can disrupt critical periods of development,¹ and one-third of children diagnosed with epilepsy between 1 and 36 months will have drug-resistant epilepsy (DRE).² Despite the importance of effective treatment, widespread treatment variability exists because epilepsy is not a single disorder but a heterogeneous group of disorders with numerous etiologies and varied natural histories.³

Most patients are initially treated with antiseizure medications (ASMs); however, after a first or second ASM fails to control seizures, the likelihood of sustained seizure freedom with any ASM substantially declines.⁴ In such contexts, other treatments including dietary therapy, surgery, or electrical stimulation devices are considered. Compared with other treatments, surgical treatments are distinctive in aiming to address underlying structural causes of epilepsies but are likely underused.^{5,6} Resection or disconnection of epileptogenic brain tissue can lead to seizure freedom (curative surgery) or seizure reduction (palliative surgery). Several factors may affect the outcomes including underlying pathology, surgery type, location, extent of resection, and concordance of presurgical evaluations. These factors affect judgements regarding epileptogenic zone identification and decisions regarding resection boundaries that aim to optimize benefits for seizure management while minimizing potential functional deficits.

Epilepsy specialists widely agree surgical treatment can be highly effective compared with serial trials of ASM in selected populations.^{7,8} Understanding current evidence for surgical treatment in early life epilepsy is critical for developing evidence-based treatment guidelines and identifying key evidence gaps. To date, systematic reviews, including a 2020 update by the National Institute for Health Care Excellence, have evaluated broader populations without focusing specifically on patients younger than 3 years of age.⁹

To assess existing evidence and characterize evidence gaps, the American Epilepsy Society (AES) identified the need for a systematic review of interventions for early life epilepsy. On behalf of the Agency of Healthcare and Quality (AHRQ) and the Patient-Centered Outcomes Research Institute (PCORI), we performed a systematic review assessing the treatment of epilepsy in children aged 1–36 months.¹⁰ In this publication, we summarize evidence on benefits and harms of surgical interventions.

Methods

Standard Protocol Approvals, Registrations, and Patient Consents

This systematic review was conducted in accordance with the AHRQ Methods Guide.¹¹ The review protocol was posted on the AHRQ website for public comment and registered in PROSPERO (CRD42021220352).

Search Strategy

To inform scope and methods, we interviewed and incorporated feedback from 9 stakeholders including neurologists, neurosurgeons, dietitians, and nurse practitioners. A professional information specialist searched EMBASE, MEDLINE, PubMed, and the Cochrane Library for studies published from January 1, 1999, to August 19, 2021 (full search strategy in the eMethods supplemental data, links.lww.com/WNL/C432).


Inclusion Criteria

Studies were required to describe outcomes among children with epilepsy undergoing surgery from 1 to 36 months of age. We excluded studies assessing treatments for infantile spasms, metabolic epilepsies, status epilepticus, and acute symptomatic seizures. Infantile spasms were excluded because of differing biology, a comparatively well-defined evidence base, and unique treatment considerations. Studies were not required to include EEG confirmation of seizures. If studies reported a mix of patient ages/seizure types, we required the study to either (1) include ≥80% relevant population or (2) report relevant data separately as a subgroup.

All study designs were considered for inclusion. Studies were required to report data for ≥10 infants per procedure. Key outcomes included seizure freedom, seizure frequency, adverse effects, all-cause mortality, sudden unexplained death in an epilepsy patient, patient quality of life (QOL), and caregiver QOL. We also extracted other outcomes including neurodevelopment and sleep quality (see Supplemental data for full list of outcomes extracted). For effectiveness outcomes, studies were required to report outcomes at ≥12 weeks. For harms outcomes, there was no minimum follow-up. Seizure freedom was defined as International League Against Epilepsy (ILAE) 1,¹² Engel Ia,¹³ or studies reporting “seizure freedom” with no further description. We also performed a sensitivity analysis to describe outcomes if seizure freedom included Engel I. For seizure frequency, favorable outcome was defined as >50% reduction in seizure frequency, Engel I or II, or ILAE I to IV.

Two analysts independently screened each abstract in DistillerSR, with disagreements resolved by consensus. For

Figure 1 Years When Surgical Procedures Were Performed



Note: Studies not appearing in this figure either were not included for effectiveness data or did not report the years when surgery was performed.

predetermined key outcomes, we rated the risk of bias using Cochrane Risk of Bias 2 for randomized controlled trials,¹⁴ the ROBINS-I instrument¹⁵ for nonrandomized studies with control groups, and Evidence-based Practice Center guidance for studies without control groups.¹⁶ For key outcomes, we also rated strength of evidence (SOE) using the 2013 AHRQ Methods Guide recommendations,¹⁷ which uses domains including study design, risk of bias, consistency of results across trials, directness, and precision.

Data Availability

Study data will be made available on reasonable request for academic purposes.

Results

Searches identified 11,123 potential citations. After title and abstract screening, 41 studies met the inclusion criteria, of which 18 studies (in 19 articles) addressed surgical interventions. No studies addressed neuromodulation. Sixteen studies described efficacy and 12 reported harms.

Effectiveness of Surgical Interventions

Sixteen prestudies/poststudies (in 17 articles) described effectiveness of surgical interventions. Although we only included studies published after 1999, surgical procedures described in these studies were performed over nearly 4 decades (Figure 1). All studies were retrospective prechart/postchart reviews except for one which used registry data.¹⁸

The number of infants meeting the inclusion criteria from each study ranged from 10 to 58. Surgical interventions were broadly categorized as hemispherectomy/hemispherotomy (anatomical hemispherectomy, functional hemispherectomy, hemispherotomy) or nonhemispheric resections (e.g., multilobar, lobar, focal resections, or disconnections). Twelve studies described infants undergoing hemispherectomy/hemispherotomy, 8 described nonhemispheric resections, and 1 study focused only on tumor resections in infants with epilepsy due to malignancy.

Five studies were conducted in the United States. The remaining 11 non-US studies were conducted in Germany¹⁹⁻²¹ (n = 3), Japan^{22,23} (n = 2), Canada^{24,25} (n = 2), Italy^{26,27} (n = 2), Sweden¹⁸ (n = 1), or included data from multiple countries (n = 1).²⁸ All US studies were single-center studies from the University of California at Los Angeles,^{29,30} University of Colorado,³¹ Cleveland Clinic,³² Boston Children's Hospital,³³ and Miami Children's hospital.³⁴ One study²⁸ included data from 19 multinational centers with surgical procedures performed from 1999 to 2020. Data from 6 patients cared for at 2 of 19 centers (University of California at Los Angeles and Cleveland Clinic) may also have been included in other studies,^{29,30,32} given the overlap in periods (author correspondence).

For many studies, data represent either subgroups or individual patient data; thus, patient characteristics such as age, seizure etiology, and the length of follow-up were variably reported. No studies reported on race.

Table 1 Seizure Freedom After Surgery, Study Characteristics

Author (Year)	Surgical procedure	Inclusion criteria	Seizure etiology or pathology	Seizure freedom	Follow-up
Hemispherectomy/Hemispherotomy					
Cook et al. 2004²⁹ Jonas et al. 2004³⁰	Anatomic hemispherectomy (n = 14) Functional hemispherectomy (n = 15) Hemispherotomy (n = 26)	Children with intractable seizures undergoing hemispherectomy at UCLA between 1986 and 2002	Cortical dysplasia	Combined: 42 of 55 (76%) Anatomical hemispherectomy: 12 of 14 (86%) Functional hemispherectomy/ hemispherotomy: 30 of 41 (73%)	2 years
Lettori et al. 2007²⁶	Anatomic hemispherectomy (n = 6) Functional hemispherectomy or hemidecortication (n = 4)	Children with drug resistant epilepsy after ≥ 3 drugs at maximal dosage treated with hemispherectomy within 5 years of age at Catholic University (Italy)	7 HME, 2 dysplastic, 1 SWS	5 of 10 (50%)	1 year
Loddenkemper et al. 2007³²	Hemispherectomy (n = 14)	50 infants younger than 3 years old among 251 consecutive pediatric patients undergoing epilepsy surgery at Cleveland Clinic between 1989 and 2001	7 HME, 5 MCD, 2 SWS	9 of 14 (64%)	Median 6 months (range 4–42) ^a
Otsuki et al. 2013²³	Hemispherotomy (n = 18)	Consecutive patients with cortical dysplasia and medically refractory epilepsy undergoing epilepsy surgery at aged younger than 6 at the National Center of Neurology and Psychiatry (Japan) from December 2000 to August 2011. Excluded patients with tuberous sclerosis, dysplastic tumors, and encephalomalacia	Cortical dysplasia	12 of 18 (66%)	NR
Pinto et al. 2014³³	Anatomic hemispherectomy (n = 10) Functional hemispherectomy (n = 4) Periinsular hemispherectomy (n = 1)	Children undergoing epilepsy surgery at Children's Hospital Boston from 1997 to 2011	Anatomic hemispherectomy patients: 6 HME, 1 polymicrogyria, 2 cortical dysplasia, 1 stroke Functional hemispherectomy/ periinsular hemispherotomy: 2 HME, 2 cortical dysplasia, 1 stroke	1 of 15 (7%)	≥ 1 year
Reinholdson et al. 2015¹⁸	Hemispherotomy (n = 12)	Swedish National Epilepsy Surgery Register data of children age younger than 4 years undergoing epilepsy surgery in Sweden between 1995 and 2010	7 HME, 2 polymicrogyria, 1 FCD I, 1 FCD unspecified, 1 gliosis/ nonspecific	7 of 12 (58%)	2 years
Roth et al. 2021²⁸	Hemispheric Procedures (n = 48)	All children undergoing epilepsy surgery at <3 months or 100 days beyond 40 weeks gestation. For inclusion, infants were required to have ≥ 6 months of follow-up, unless the patient died	NR for this subgroup	70% (extrapolated to be 30 of 43) ^a	Median 51 months (IQR 27 to 126)
Schramm et al. 2012²⁰	Hemispherotomy (n = 21)	All patients younger than 18 undergoing hemispherotomy at the Bonn University Medical Center (Germany) operated on between 1990 and the end of 2009 with ≥ 1 year follow-up.	NR for this subgroup	16 of 21 (76%)	≥ 1 year

Continued

Table 1 Seizure Freedom After Surgery, Study Characteristics (*continued*)


Author (Year)	Surgical procedure	Inclusion criteria	Seizure etiology or pathology	Seizure freedom	Follow-up
Multilobar, Lobar, or Focal Resection					
Loddenkemper et al. 2007³²	Focal cortical resection (n = 10)	50 infants younger than 3 years old among 251 consecutive pediatric patients (younger than 18 years old) undergoing epilepsy surgery at Cleveland Clinic between 1989 and 2001 were considered for inclusion	7 malformation of cortical development without HME (MCD), 2 MCD + ganglioma, 1 TS	7 of 10 (70%)	Median 6 months (range 4–42) ^b
Reinholdson et al. 2015¹⁸	Frontal lobe resection (n = 12) Temporal lobe resection (n = 12)	Swedish National Epilepsy Surgery Register data of children aged younger than 4 undergoing resective epilepsy surgery in Sweden between 1995 and 2010	Temporal lobe resection: 2 focal cortical dysplasia II, 3 FCD unspecified, 2 gliosis/nonspecific, 4 low grade tumor, 1 TS Frontal lobe resection: 2 FCD II, 5 FCD unspecified, 2 gliosis or nonspecific, 2 TS, 1 vascular malformation	Combined: 12 of 24 (50%) Temporal lobe resection: 8 of 12 (67%) Frontal lobe resection: 4 of 12 (33%)	2 years
Sugitomo et al. 1999²⁴	Focal cortical resection (n = 10)	Children, aged 0–3 years, who had epilepsy surgery at The Hospital for Sick Children, Toronto (Canada) from 1991 to 1996	NR	4 of 10 (40%)	Mean 3.2 years (range 0.25–6.7 years)
Kalbhenn et al. 2019²¹	Posterior disconnection (n = 10)	Consecutive patients undergoing posterior disconnection surgery between 2005 and 2017 for refractory posterior quadrant epilepsy at Evangelisches Klinikum Bethel (Germany)	6 FCD, 2 ganglioma + FCD, 1 polymicrogyria, 1 meningeal angiomas	5 of 10 (50%)	2 years
Roth et al. 2021²⁸	Focal resections (n = 16)	All children undergoing epilepsy surgery at <3 months or 100 days beyond 40 weeks gestation. For inclusion, infants were required to have ≥6 months of follow-up, unless the patient died	NR for this subgroup	56% (extrapolated to be 9 of 16)	Median 24 months (5–55 IQR)

FCD: Focal cortical dysplasia; HME: Hemimegalencephaly; MCD: Malformation of cortical development; NR: Not reported; TS: Tuberous sclerosis; SWS: Sturge-Weber syndrome.

^a N provided by author correspondence.

^b Follow-up duration for hemispherectomy and focal cortical resections combined.

Figure 2 Seizure Freedom Rates after Hemispherectomy or Hemispherotomy



The vertical bars display 95% confidence intervals. When multiple studies reported the same follow-up times, we used horizontal offsets to improve visibility. Only 1 study (Cook) reported data across multiple time points. For studies only reporting a minimum follow-up interval (e.g., follow-up >1 year after surgery), data were plotted at that minimum time point. Of note, Otsuki et al. was not included as the study did not report follow-up duration for this subgroup. 95% confidence intervals are shown for each data point. All studies were prestudies/poststudies, so the precise cause of seizure freedom cannot be easily determined.

Hemispherectomy/Hemispherotomy

Seizure Freedom

Eight retrospective prestudies/poststudies (in 9 articles) included 188 infants^{18,20,23,26,28-30,32,33} and reported on seizure freedom from 6 months to mean 4.3 years after surgery (Table 1). One study²³ did not report the follow-up interval for the subgroup of included patients. All studies were assessed as a high risk of bias for many reasons including lack of a control group.

Seizure freedom rates at 1 year ranged from 7% to 76% (Figure 2). Six studies described outcomes at ≥ 1 year. Two larger studies reported seizure freedom rates of 76% (42/55)²⁹ and 70% (30/43)²⁸ at 2 years and median 4.3 years, respectively. Three smaller studies reported rates ranging from 50% to 76%.^{18,20,26} Finally, 1 study reported only 7% (1/15) infants were seizure free at follow-up of ≥ 1 year after surgery.³³

We used a stringent definition of seizure freedom, which required studies using the Engel classification to report Engel Ia. However, if we had considered Engel I as seizure freedom, seizure freedom in Pinto et al.³³ would increase to 66% (10 of 15); also, 4 other studies^{19,22,25,31} reporting rates of 55%–81% (consistent with the range of seizure freedom rates we already identified) would have been included.

Notably, prestudies/poststudies lack true control groups, posing a challenge for knowing what would have happened had these patients not undergone surgery. However, most infants underwent surgery for intractable epilepsy, suggesting none would have experienced seizure freedom without surgery. In addition, although seizure counts were assessed using retrospective data from charts (i.e., not captured in the

context of a trial), seizure freedom is less subject to recall bias or other types of bias than other outcomes such as seizure frequency. Thus, we concluded that low strength evidence suggests some infants with intractable epilepsy achieve seizure freedom after hemispherectomy/hemispherotomy (Table 2).

Seizure Frequency

Nine retrospective prestudies/poststudies (including 186 infants) reported on seizure frequency. All studies reported more than half of infants achieved a favorable outcome (Engel I or II, ILAE I to IV, or >50% seizure reduction). The proportion of infants achieving favorable outcome ranged from 67% to 100%, with most studies reporting a follow-up of at least 1 year. Specifically, studies reported the following proportion of infants had favorable outcomes at follow-up: 67% (10/15),³³ 72% (13/18),²³ 73% (35/48),²⁵ 72% (31/43),²⁸ 80% (8/10),²² 88% (14/16),³¹ 92% (11/12),¹⁸ 93% (13/14),³² and 100% (10/10).²⁶ However, seizure counts were assessed using retrospective data from charts (i.e., not captured in the context of a trial), which we assessed as high risk of bias. Thus, the evidence was considered insufficient to draw a conclusion regarding seizure frequency.

Favorable outcome after epilepsy surgery could depend on multiple factors, including underlying etiology. Eight studies^{21,22,24,26,31,32,34} reported individual patient data for pathology or etiology, surgical intervention, and outcomes including 65 infants undergoing hemispherectomy or hemispherotomy. Seizure etiology or pathology was reported as hemimegalencephaly (HME) (58%), focal cortical dysplasia (FCD) or malformation of cortical development (MCD) without HME (20%), or other pathology (22%). A summative analysis found the proportion of infants achieving favorable outcomes was similar across these 3 groups: 89% (34/38)

Table 2 Strength of Evidence for Surgical Procedures: Efficacy and Harms

Treatment	Outcome	Study findings	Strength of evidence	Conclusion
Hemispherectomy/ Hemispherotomy	Seizure Freedom	Seizure freedom rates ranged from 7% to 76% at 1 year (8 prestudies/poststudies [in 9 articles], ^{18,20,23,26,28-30,32,33} combined 188 infants)	Low	Some infants with medically refractory epilepsy may achieve seizure freedom after hemispherectomy/hemispherotomy.
	Favorable outcome (>50% seizure reduction)	The proportion of infants achieving a favorable outcome ranged from 67% to 100%. (9 prestudies/poststudies, ^{18,22,23,25,26,28,31-33} combined 186 infants)	Insufficient	Evidence is insufficient because of study limitations.
Anatomic hemispherectomy	Surgical mortality	A single death ²⁹ was reported across 3 studies. (3 prestudies/poststudies, ^{28,29,35} combined 26 infants)	Insufficient	Evidence is insufficient because of sparse data.
	Hydrocephalus	Rates of hydrocephalus ranged from 43% to 100% (3 prestudies/poststudies, ^{26,33,35} combined 19 infants)	Insufficient	Evidence is insufficient because of sparse data.
Functional hemispherectomy/ hemispherotomy	Surgical Mortality	A single death ³¹ was reported across 8 studies. (8 prestudies/poststudies, ^{19,22,23,28,29,35,36} combined 180 infants)	Low	Surgical mortality after functional hemispherectomy/hemispherotomy is rare.
	Hydrocephalus	Rates of hydrocephalus ranged from 0% to 33%. (9 prestudies/poststudies, ^{18,19,20,25,26,31,33,35,36} combined over 96 ^a)	Insufficient	Evidence is insufficient because of study limitations.
Nonhemispheric procedures (multilobar, lobar, or focal cortical resection)	Seizure Freedom	Rates of seizure freedom ranged from 40% to 70% (5 prestudies/poststudies, ^{18,21,24,28,32} combined 70 infants)	Low	Some infants with medically refractory epilepsy may achieve seizure freedom after intralobar, multilobar, or focal cortical resection.
	Favorable outcome (>50% seizure reduction)	The proportion of infants achieving a favorable outcome ranged from 50% to 85%. (6 prestudies/poststudies), ^{18,24,25,28,32,34} 131 combined infants)	Insufficient	Evidence is insufficient because of study limitations.
Multilobar, intralobar, or focal resections	Surgical Mortality	No deaths were reported. (3 prestudies/poststudies, ^{28,35,36} combined 82 infants)	Low	With multilobar/lobar/focal resection, surgical mortality is rare.
	Hydrocephalus	3 cases of postoperative hydrocephalus were reported. (5 prestudies/poststudies, ^{19,25,28,35,36} combined over 108 infants)	Low	Hydrocephalus requiring shunt placement after multilobar, unilobar, or focal resections is uncommon

^a One study¹⁹ only reported a percentage and thus was not included in this number.

for HME, 92% (12/13) for FCD/MCD without HME, and 93% (13/14) for other pathology.

Nonhemispheric Surgical Procedures

Seizure Freedom

Five prestudies/poststudies including 70 infants reported on seizure freedom. Infants underwent focal cortical resections in 3 studies,^{24,28,32} frontal or temporal lobe resection in 1 study,¹⁸ and posterior disconnection in 1 study.²¹ The rates of seizure freedom ranged from 40% to 70%. Figure 3 presents seizure freedom rates and follow-up durations. Specifically, focal resection was followed by seizure freedom in 70% (7/10),³² 56% (9/16),²⁸ and 40% (4/10)²⁴ at a median 6, 24 months, and mean 3.2 years, respectively. Kalbhenn et al.²¹ reported that 50% (5/10) of patients were seizure free after posterior disconnection surgery at 2 years after surgery. Reinholdson et al.¹⁸ reported 50% (12/24) of children undergoing frontal or temporal lobe resection were seizure free at 2 years after surgery. If seizure freedom had included studies reporting Engel I, 3 additional studies reporting the


rates of 62%,¹⁹ 69%,³⁴ and 91%²⁵ would have also been included.

We concluded some infants with intractable epilepsy achieve seizure freedom after intralobar, multilobar, or focal cortical resection (SOE: Low).

Seizure Frequency

Seven retrospective prestudies/poststudies including 148 infants reported seizure frequency. Six of 7 studies reported seizure frequency data that allowed for determination of favorable outcome for seizure frequency.^{18,24,25,28,32,34} All studies found more than half of infants achieved a favorable outcome for seizure frequency: specifically, the proportion of infants achieving favorable outcomes was 50% (5/10),²⁴ 83% (20/24),¹⁸ 85% (11/13),³⁴ 90% (52/58),²⁵ 94% (15/16),²⁸ and 100% (10/10).³² The mean follow-up was ≥1 year after surgery for all studies. A seventh study¹⁹ reported of 17 infants for whom the extent of final resection was intralobar, 76% (13/17) were Engel I, whereas 24% (4/17) were Engel II

Figure 3 Seizure Freedom Rates after Multilobar, Lobar, or Focal Resection



to IV. Nevertheless, because of a high risk of bias, evidence was judged insufficient to draw a conclusion.

Studies reporting individual patient data included 43 infants undergoing multilobar, lobar, or focal resection. Seizure etiology or pathology was FCD/MCD without HME (56%) or other pathologies (44%). A simple summative analysis found the proportion of infants with favorable outcome rates was 67% (16/24) for FCD/MCD without HME and 74% (14/19) for other pathologies.

Tumor Resection

One retrospective chart review²⁷ focused exclusively on 20 infants with epilepsy because of primary supratentorial brain tumors. Mean time from tumor diagnosis to surgery was 0.86 (SD 0.63) months. Of 17 patients with 8 years postoperative follow-up, 53% (9/17) were Engel I, 24% (4/17) Engel II, 12% (2/17) Engel III, and 12% (2/17) Engel IV.

Developmental Outcomes (All Procedures)

Only 4 prestudies/poststudies reported on developmental outcomes including developmental quotient [DQ], language, or functional status. Three studies^{26,30,32} focused on hemispherectomy, including 2 studies reporting DQ after hemispherectomy. Loddenkemper et al.³² included 24 infants undergoing hemispherectomy or focal resection at a median age of 14 months (3–34). Infants were evaluated at a median 12 months (3–34) preoperatively and at a median 24 months (10–53) after surgery. The proportion of infants with developmental delay (DQ < 70) decreased after surgery, but was not statistically significant ($p = 0.125$). Furthermore, 52% (26/50) of consecutive infants were excluded because of incomplete data or the use of other neuropsychological tests, potentially limiting generalizability. Jonas et al.³⁰ found that in 16 infants undergoing hemispherectomy for HME, at 6–24 months after surgery, the Vineland DQ increased by 9.1 (SD 16) compared with presurgery. The spoken language rank also increased from 0.33 (SD 0.5) to 1.4 (SD 1.8) after

surgery. Lettori²⁶ included 10 infants meeting the inclusion criteria and undergoing hemispherectomy. Before surgery, 20% (2/10) had a dependent functional status, and functional status could not be assessed for 80% (8/10). After surgery, 6 were dependent, 3 semiindependent, and 1 independent.

Finally, one study reported some infants had improvement in developmental delay after undergoing focal cortical resection (8/10 with delay preoperatively, 6 infants with improved/good status after surgery).²⁴ However, the study did not report how delay was assessed.

Harms

Twelve prestudies/poststudies described harms after surgery (Table 3).

Hemispherectomy/Hemispherotomy

Eleven prestudies/poststudies^{18,19,22,23,25,26,28,29,31,33,35} reported harms after hemispherectomy/hemispherotomy. Roth et al.²⁸ included data from 19 centers with surgical procedures performed from 1999 to 2020. Seven patients cared for at 3 of 19 centers (University of California at Los Angeles, Cleveland Clinic, and Great Ormond Street Hospital) may also have been previously reported in other studies included in this report (author correspondence).^{29,32,35} Iwasaki et al.³⁶ described harms for 75 infants, of which 9 hemispherectomy patients had previously been described in another study,²³ also included in this report (author correspondence).

Mortality

Nine prestudies/poststudies reported on mortality after hemispherectomy/hemispherotomy. For the outcome of surgical mortality, studies were assessed as low risk of bias. Studies described mortality after anatomical hemispherectomy (3 studies), functional hemispherectomy or hemispherotomy (8 studies), or across multiple procedures (lesionectomy, cortical resection, and hemispherectomy/hemispherotomy, 1 study).

Table 3 Studies Reporting on Harms

Author (Year)	Surgical procedure (n)	Inclusion criteria	Seizure etiology or pathology	Harms reported (follow-up interval)
Cook et al. 2004 ²⁹	Anatomical hemispherectomy (n = 14) Functional hemispherectomy (n = 15) Hemispherotomy (n = 26)	Children with intractable seizures undergoing hemispherectomy at UCLA between 1986 and 2002	Cortical dysplasia	Mortality (NR) Other AEs
Dunkley et al. 2010 ³⁵	Anatomical hemispherectomy (n = 2) Functional hemispherectomy (n = 25) Multilobar/Lobar/Focal Resection (n = 15)	All children undergoing resection for epilepsy at younger than 36 months of age and followed up for at least 2 years at Great Ormond Street Hospital	NR by procedure	Mortality (NR) VP or LP shunt placement (placed at 6 weeks, 6 months, 12 months, 4 years)
Iwasaki et al. 2015 ²²	Hemispherotomy (n = 10)	Consecutive patients undergoing hemispherotomy for treatment of intractable epilepsy in Tohoku University Graduate School of Medicine between 2001 and 2012	NR	Mortality (NR) VP shunt (NR)
Kadish et al. 2019 ¹⁹	Hemispherotomy (n = 22)	Consecutive patients undergoing presurgical evaluation at Epilepsy Center Freiburg (Germany) with follow-up > 1 year after surgery	NR by procedure	Mortality ("Perioperative") VP shunt (NR) Other AEs
Kumar et al. 2015 ³¹	Hemispherotomy (n = 16)	All children under the age of 1 year undergoing surgical intervention to treat medically refractory epilepsy at Children's Hospital of Colorado between 2002 and 2013	NR	Mortality ("Perioperative") Hydrocephalus (mean follow-up for study 56 months, range 3–133 months) Other AEs
Lettori et al. 2007 ²⁶	Anatomic hemispherectomy (n = 6) Functional hemispherectomy or hemidecortication (n = 4)	Patients treated with hemispherectomy within 5 years of age in the Child Neurosurgery Unit of Catholic University, Rome from 1980 to December 2003, "we enrolled in the study only 19 thoroughly studied children, drug resistant with at least 3 drugs at maximal dosage with no seizure control"	7 HME, 2 dysplastic, 1 SWS	Hydrocephalus (NR) Other AEs
Otsuki et al. 2013 ²³	Hemispherotomy (n = 18)	Consecutive patients with cortical dysplasia and medically refractory epilepsy undergoing epilepsy surgery at age <6 at the National Center of Neurology and Psychiatry (Japan) from December 2000 to August 2011. Excluded patients with tuberous sclerosis, dysplastic tumors, and encephalomalacia	Cortical dysplasia	Mortality (NR)
Pinto et al. 2014 ³³	Anatomic hemispherectomy (n = 10), Functional hemispherectomy (n = 4), Peri-insular hemispherectomy (n = 1)	Children undergoing epilepsy surgery (anatomic hemispherectomy, functional hemispherectomy, and periinsular hemispherotomy) at Children's Hospital Boston from 1997 to 2011. Excluded patients with progressive disease including Rasmussen encephalitis and Sturge-Weber syndrome	Anatomic hemispherectomy patients: 6 HME, 1 polymicrogyria, 2 cortical dysplasia, 1 stroke Functional hemispherectomy/Periinsular hemispherotomy: 2 HME, 2 cortical dysplasia, 1 stroke	VP shunt (NR)
Reinholdson et al. ¹⁸	Hemispherotomy Frontal lobe resection Temporal lobe resection	Swedish National Epilepsy Surgery Register data capturing population based, observational cohort of children under 4 years of age undergoing resective epilepsy surgery in Sweden between 1995 and 2010	7 HME, 2 polymicrogyria, 1 FCD I, 1 FCD unspecified, 1 gliosis/nonspecific	VP shunt (within 2 years after surgery)

Continued

Table 3 Studies Reporting on Harms (*continued*)

Author (Year)	Surgical procedure (n)	Inclusion criteria	Seizure etiology or pathology	Harms reported (follow-up interval)
Roth et al. 2021²⁸	Hemispheric surgeries (25 peri-insular, 12 vertical functional hemispherectomies, 10 anatomical hemispherectomies, 1 unknown) Focal resections (7 lobectomies, 12 lesionectomy)	Epilepsy surgery at <3 months age or 100 days beyond 40 weeks gestation at one of 19 centers. For inclusion, infants were required to have ≥6 months of follow-up, unless the patient died. Excluded infants with epilepsy because of hypoxic ischemic encephalopathy.	28 cortical dysplasia, 17 hemimegalencephaly, 5 tubers, 4 nonspecific findings, 1 glioneuronal hamartoma, 1 stroke, 1 Sturge Weber, 1 hematoma, 6 unknown	Mortality (“Perioperative”) Hydrocephalus (median follow-up for all study outcomes 51 months, IQR 21 to 126) Other AEs (combined incidence across all procedures)
Iwasaki et al. 2021³⁶	Hemispherotomy (periinsular hemispherotomy, vertical parasagittal hemispherotomy) (n = 27) Multilobar surgery, Unilobar surgery (n = 48)	Infants undergoing “first curative epilepsy surgery” at <3 between August 2006 and February 2019 at the National Center of Neurology and Psychiatry, Tokyo, Japan ≥1 year follow-up postoperative, and post-operative developmental assessment at 1 year or last follow-up after 2 years or longer. Excluded patients who had undergone palliative procedures (corpus callosotomy and vagus nerve stimulation implantation).	22 hemimegalencephaly, 33 other malformations of cortical development, 10 low-grade developmental tumors, 6 tuberous sclerosis complex, 3 Surge-Weber syndrome, 1 perinatal ischemia.	Mortality (patients required to have ≥1 year follow-up after surgery) Hydrocephalus requiring surgical intervention (NR)
Steinbok et al. 2009²⁵	Lesionectomy, cortical resection, and hemispheric surgery (n = 151 procedures in 116 patients)	Age <3 years undergoing epilepsy surgery at multiple centers across Canada from January 1987 to September 2005. Patients undergoing surgery for a lesion, such as a tumor, who happened to present with seizures but for whom the surgery was performed for the lesion rather than the epilepsy were excluded.	NR by procedure	Mortality (intraoperative) Hydrocephalus (within a “few months” after surgery) Other AEs

Abbreviations: FCD = focal cortical dysplasia; HME = hemimegalencephaly.

Nine infants reported in Iwasaki 2021³⁶ had previously been reported in Otsuki et al. 2013. For Steinbok et al.²⁵ the number of hemispheric vs nonhemispheric procedures for which hydrocephalus was reported was unclear.

Only 3 retrospective chart reviews reported data for infants undergoing anatomical hemispherectomy. Cook et al.²⁹ reported of 14 infants: there was 1 intraoperative death, an 8-month-old infant with HME. Two additional studies with 12 infants^{28,35} reported no deaths. Evidence was insufficient to permit conclusions because of sparse data.

Eight studies described surgical mortality for a combined 196 infants undergoing functional hemispherectomy/hemispherotomy. Seven of 8 studies (including 180 infants)^{19,22,23,28,29,35,36} reported no deaths. Kumar et al.³¹ reported of 16 hemispherotomies, 1 death occurred in an infant with epidermal nevus syndrome, right HME, and multiple other congenital abnormalities who developed refractory seizures after surgery and technological support was withdrawn.

Steinbok et al.²⁵ reported a single death across 116 infants undergoing 151 surgical procedures (hemispherectomy/hemispherotomy, lesionectomy, or cortical resections). The intraoperative death occurred in a 3.9 months child with tuberous sclerosis undergoing attempted resection of intraventricular and extraventricular lesions.

Overall, these studies suggest perioperative mortality after functional hemispherectomy or hemispherotomy is uncommon. However, these studies were primarily single center retrospective chart reviews including heterogeneous infants (with many different seizure etiologies), and studies often failed to specify the proportion of infants not included because of missing data. Furthermore, it is possible that centers with higher mortality rates might choose not to publish their data. However, despite these study limitations, we concluded surgical mortality after functional hemispherectomy/hemispherotomy is rare (SOE: Low).

Hydrocephalus

For anatomical hemispherectomy, 3 studies (combined 19 surgeries) reported hydrocephalus and/or need for a ventriculoperitoneal shunt (VPS) were common. Dunkley et al.³⁵ reported 2/2 infants undergoing anatomical hemispherectomy required VP shunt placement 12 months after surgery. Similarly, Pinto et al.³³ reported 7/10 infants undergoing anatomical hemispherectomy required VP shunt placement (follow-up interval NR). Lettori et al.²⁶ reported 3/7 infants undergoing anatomical hemispherectomy or hemidecortication developed hydrocephalus (follow-up interval NR).

For functional hemispherectomy or hemispherotomy, 9 studies (combined n = 96, plus infants from 1 study¹⁹ only reporting a percentage, and another study²⁵ with an unclear denominator) reported on hydrocephalus. Studies reported lower rates of hydrocephalus/VPS compared with anatomical hemispherectomy. One of 9 studies reported no infants (0/10) developed hydrocephalus.²² Another study reported 4 infants undergoing functional hemispherectomy developed hydrocephalus within a few months after surgery; at least 22 infants underwent functional hemispherotomy in this study,

but the total number of infants undergoing this procedure was unclear.²⁵ The remaining 7 studies reported rates of 8% (1/12),¹⁸ 11% (3/27),³⁵ 16% (n NR),¹⁹ 20% (1/5),³³ 22% (6/27),³⁶ 25% (4/16),³¹ and 33% (1/3).²⁶

Finally, Roth et al.²⁸ reported hydrocephalus in 25% (11 of 44) infants undergoing either anatomical hemispherectomy or functional hemispherectomy/hemispherotomy. Notably, only 1 study³⁵ reported when hydrocephalus occurred, although a second study²⁵ reported a time range. Four studies^{19,22,26,36} did not report when hydrocephalus occurred, and the remaining studies provided a time point at which other outcomes were measured (e.g., >1 year after surgery) but no other information regarding the timing of hydrocephalus.

Given multiple factors including heterogeneity across patients and procedures and inconsistent outcome reporting, evidence was deemed insufficient to draw a conclusion regarding hydrocephalus and/or VPS after hemispherectomy/hemispherotomy.

Multilobar, Lobar, and Focal Resections

Mortality

Four prestudies/poststudies described surgical mortality for infants undergoing nonhemispheric procedures. Three studies^{28,35,36} described surgical mortality for a combined 82 infants undergoing multilobar, lobar, or focal resections and reported no deaths. These 3 studies included infants undergoing a range of nonhemispheric procedures. Dunkley et al.³⁵ included 15 infants undergoing either multilobar, lobar, or focal resections. Iwasaki et al. 2021³⁶ included 48 infants undergoing multilobar (13 posterior quadrantic disconnections, 5 multifocal cortical resections, 1 subtotal hemispherotomy) or unilobar surgeries (16 focal cortical resections or lesionectomies, 8 anterior temporal lobectomies, and 5 frontal lobectomies or disconnections). Roth et al.²⁸ included 19 infants undergoing focal resections.

A fourth study, Steinbok et al.²⁵ reported only a single mortality across 116 infants undergoing 151 procedures, which were either a hemispherectomy/hemispherotomy, lesionectomy, or cortical resections.

This evidence base for mortality after nonhemispheric procedures is small with important limitations. All studies were retrospective chart reviews, and 2 studies reported experience drawn from single centers. However, the results from Roth et al.¹⁸ (which included data from 19 centers) were consistent in also reporting no deaths. Reported mortality rates may be artificially low if centers with higher mortality rates choose not to publish their data. Nevertheless, despite these limitations, we concluded surgical mortality after multilobar, lobar, or focal resection is rare (SOE: Low).

Hydrocephalus

Four studies^{19,28,35,36} with a combined 108 procedures reported on infants undergoing focal, intralobar, or multilobar resections. No patients developed hydrocephalus (follow-up duration NR for 2 studies,^{19,35} median of 24 months,²⁸ and >1

year after surgery³⁶). Despite study limitations, we concluded that hydrocephalus after multilobar, intralobar, or focal surgery is rare (SOE: Low).

Additional Adverse Effects (All Procedures)

Reporting of other AEs was inconsistent. Two studies^{26,29} reported other AEs after anatomic hemispherectomy including infection, transient fever, cranial nerve III palsy, subdural fluid collection, and CSF leakage. For functional hemispherectomy/hemispherotomy, in addition to infection,^{26,29} AEs reported included intraoperative disseminated intravascular coagulation (1/37),²⁸ acute postsurgical seizures (23%),¹⁹ epidural hemorrhage requiring surgical revision (1/22),¹⁹ dural adhesions requiring late reoperation (1/41),²⁹ pituitary failure because of thalamic lesion (1/22),¹⁹ cerebral salt wasting syndrome (2/27),³⁶ diabetes insipidus (3/27),³⁶ sinus thrombosis resulting from diabetes insipidus (2/27),³⁶ and asymptomatic hemorrhagic infarction (1/27).³⁶

For multilobar, lobar, or focal resections, 1 study reported 1/10 infants developed transient hemiparesis after posterior disconnection for refractory posterior quadrant epilepsy.²¹ Another study³⁶ reported the following complications requiring surgical or medical intervention in 48 infants undergoing multilobar, unilobar, or focal resections: subdural hygroma (n = 3), cyst formation (n = 2), asymptomatic cerebral infarction (n = 1), bacterial meningitis (n = 1), and psychiatric symptoms (n = 1).

Discussion

Our findings suggest surgical interventions for children aged 1–36 months with epilepsy can be beneficial for reducing seizures for some children, and surgical mortality is rare. Although other outcomes including developmental/cognitive outcomes, sleep, and QOL are also important, few or no studies reported these. Overall, the evidence base remains sparse, with key limitations, including a lack of prospective controlled studies, and inadequate measurement of important outcomes. Despite including studies reporting as few as 10 patients per procedure, we identified only 18 studies, of which all were prestudies/poststudies and 17 were retrospective. Notably, the absence of rigorous trials in this age group does not demonstrate that surgery is ineffective, instead it highlights a critical evidence gap.

Despite these limitations, the rates of seizure freedom for infants undergoing hemispherectomy/hemispherotomy or other resections (multilobar, lobar, or focal resections) reported in studies were encouraging. For hemispherectomy/hemispherotomy, with 1 exception, studies reported more than half of infants were seizure free at follow-up. Similarly, for multilobar/lobar/focal resections, studies reported seizure freedom rates of 40%–70%. Furthermore, we found low strength evidence suggesting surgical mortality was rare. As seizure freedom rates with medical management in children 1–36 months old with DRE are substantially lower than 40%–70%, these findings suggest epilepsy surgery can be beneficial for treating seizures in this age group.

Some studies reported on surgeries performed 4 decades ago, raising questions regarding generalizability, given the changes in clinical care over time. However, most studies reported seizure freedom rates similar to those reported by Roth²⁸ (56% and 70% for focal and hemispheric procedures, respectively) a larger recent study which included patients from multiple centers operated on from 1999 to 2020. A recent study (published subsequent to our search dates) reported of 34 children <3 undergoing epilepsy surgery since 2018, 59% were Engel I outcome at median follow-up of 21.9 months.³⁷

Several factors may limit applicability of these findings. Ideally, outcomes after surgery would be reported by etiology, given the wide range of etiologies with unique clinical considerations and trajectories that may lead to DRE and evaluation for epilepsy surgery. However, sparsity of studies, clinical heterogeneity of included patients, and limitations of study reporting precluded this type of assessment. Limited reporting of clinical details also precluded consideration of other clinical factors (e.g., the number of ASM at outcome reporting, number of previous surgeries, variation in surgical procedures) on outcomes.

Most studies were small and single center, reflecting outcomes from single epilepsy surgery programs and/or surgeons. Furthermore, nearly all studies were retrospective chart reviews at risk for inconsistencies in data collection and reporting. One study of consecutive infants included only <50% (24/50) of potentially eligible infants because of missing data,³² illustrating the potential for bias from studies using a retrospective chart review design. However, few studies reported the proportion of patients excluded because of missing data. Although only studies published after 1999 were included, surgeries described were performed over 4 decades. Although excluding studies published before 1999 could have excluded relevant data, including older studies could also have resulted in inclusion of even older and potentially less generalizable data.

Although we found existing evidence to be sparse and low quality, notably, the lack of high quality studies does *not* demonstrate that surgical treatments are ineffective. Instead, it highlights the need for additional higher quality evidence. The scope of our review was limited to children aged 1–36 months because of AES's request to focus on this particular population, feasibility considerations, and resources. Although exploration of indirect evidence (e.g., studies performed in older children) could provide useful information, this was not feasible given resource constraints.

To improve the SOE, improvements to study design and data reporting are needed. In 2012, an Institute of Medicine report named long-term prospective studies assessing effects of epilepsy surgery on cognitive function with inclusion of appropriate control groups as a research priority.³⁸ Our findings demonstrate this remains an evidence gap for surgical treatments in early life epilepsy. Prospective studies with clear and consistent reporting of variables including seizure etiology,

semiology, previous and concomitant treatments, and follow-up interval are needed. As others have noted, seizure freedom remains challenging to define and key differences between the Engel classification and ILAE outcome scales pose challenges for comparing results from studies using these scales.³⁹

Pragmatic and ethical concerns exist regarding randomizing infants with epilepsy to surgery vs sham or placebo. However, a feasible next step would be a high-quality prospective, multicenter observational cohort study. This could be facilitated by a multicenter registry with standardized measures (including developmental outcomes, QOL outcomes, and adverse effects). This type of registry would offer important advantages: 1) given the relatively small number of infants undergoing surgical interventions, gathering data across multiple centers would increase the ability to measure efficacy/harms and avoid potential duplicate reporting of patients in studies, 2) improve generalizability by minimizing differences specific to individual institutions or surgeons, 3) facilitate consensus about outcome measurement (including key outcomes and follow-up duration), and 4) provide a framework for prospective efficient collection of standardized data.⁴⁰ Existing consortiums could play a role in facilitating development.

Development of core outcomes specific to patients with early life epilepsy could also support these efforts. Outcomes identified as important by stakeholders interviewed during protocol development including seizure freedom, seizure frequency, seizure severity, Engel classification, all-cause mortality, hospitalization, neurodevelopmental outcomes, QOL, sleep quality, caregiver QOL, treatment cost, and other adverse events. Given the range of seizure etiologies and surgical interventions, future studies should not only report these outcomes but also report outcomes separately for different seizure etiologies (i.e., structural vs acquired) and surgeries (i.e., focal cortical resection vs frontal lobectomy). Even some structural lesions may be further divided by detailed pathologic assessments or genetic etiologies. Without this information, future systematic reviews are likely to encounter similar difficulty drawing conclusions about specific etiologies or procedures in this age group.

Acknowledgment

Mention of individuals in this section does not imply endorsement of any part of the article; the methods and conclusions are those of the authors alone. The authors gratefully acknowledge the following individuals for their expert guidance: Adam Hartman, MD (NINDS), David Niebuhr, MD, MPH, MSc (AHRQ), Jennie Dalton, MPH (PCORI), and Joy Keller, MS, RD, MSLIS (AES). We also acknowledge support from the following individuals at ECRI: Janice Kaczmarek, Lindsey Miller, PT, DPT, and Jacki Hostetter (project management), Laura Koepfler MLS (searches), Helen Dunn and Katherine Donahue (references and article procurement), and Jennifer Maslin (formatting). The authors appreciate the input of various societies and individuals at different stages of this review: American

Epilepsy Society (submitted PCORI topic nomination, advised on topic development, served as nonsponsoring partner through AHRQ planning, and provided input on early drafts of the full report); An early draft of the full report was reviewed but not endorsed by the American Association of Neurological Surgeons/Congress of Neurological Surgeons Section on Pediatric Neurological Surgery; Emily Spelbrink (Stanford University, on behalf of the Pediatric Epilepsy Research Consortium Early Life Epilepsies Special Interest Group) provided input on an early draft of the full report; Anne Berg (Northwestern University) served on the Technical Expert Panel and provided input on an early draft of the full report; Robyn Blackford (Ann & Robert H. Lurie Children's Hospital of Chicago) served as a peer reviewer on the draft full report; Jeffrey Blount (University of Alabama at Birmingham) served as a peer reviewer on the draft full report; Kevin Chapman (Phoenix Children's Hospital) served on the Technical Expert Panel and provided input on an early draft of the full report; Jennifer Coffman (Children's Hospital Colorado) served as a peer reviewer on the draft full report; Erin Fecske (Children's Mercy Hospital) served on the Technical Expert Panel and provided input on an early draft of the full report; William Davis Gaillard (Children's National Hospital) provided input at early stages of topic formulation and provided input on an early draft of the full report; Zachary Grinspan (New York-Presbyterian Kormansky Children's Hospital) served as a peer reviewer on the draft full report; Mary Anne Meskis (Dravet Syndrome Foundation) provided input on an early draft of the full report; Gary W. Matthern (UCLA) provided input on an early draft of the full report; Solomon Moshe (Albert Einstein College of Medicine) provided input on an early draft of the full report; Douglas Nordli (University of Chicago Medicine) served on the technical expert panel and provided input on an early draft of the full report; Edward (Rusty) Novotny (Seattle Children's Hospital Center for Integrative Brain Research, University of Washington) served as a peer reviewer on the draft full report; Chima Oluigbo (Children's National Hospital) served on the technical expert panel and provided input on an early draft of the full report; Phillip Pearl (Children's Hospital Boston) provided input on an early draft of the full report; Heidi Pfeifer (Boston Children's Hospital) served on the technical expert panel; Leah Schust Myers (FamilieSCN2A Foundation) served as a key informant to determine scope; Renee Shellhaas (University of Michigan) served as a key informant to determine scope; Shlomo Shinnar (Albert Einstein College of Medicine) provided input at early stages of topic formulation and provided input on an early draft of the full report; Ann Soddors provided input on an early draft of the full report; Howard Weiner (Texas Children's Hospital and Baylor College of Medicine) served on the technical expert panel and provided input on an early draft of the full report; M.A. Whelan provided input on an early draft of the full report.

Study Funding

This work was supported by Contract No. 75Q80120D00002 from the Agency for Healthcare Research and Quality, U.S.

Department of Health and Human Services, and by the Patient-Centered Outcomes Research Institute (PCORI) through a memorandum of Agreement Amendment, number 20-603M-19.

Disclosure

A.Y. Tsou, M.J. Wu, and J.R. Treadwell report no disclosures; SK Kessler has served as an investigator for Esai, Neurocrine Biosciences, SK Life, and UCB Pharma (for which funding has been granted to Children’s Hospital of Philadelphia); she serves as a consultant and executive committee member of the Epilepsy Study Consortium (funding to Children’s Hospital of Philadelphia) and has received fees from American Academy of Neurology for educational activities; NS Abend has received funding from NIH (NINDS) K02NS096058 (to institution), Wolfson Foundation (to institution), PCORI (to institution), UCB Pharma (to institution), Epilepsy Foundation (consulting), and Demos Publishing (royalties); S.L. Massey has no disclosures. Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures.

Publication History

Received by *Neurology* November 22, 2021. Accepted in final form June 9, 2022. Submitted and externally peer reviewed. The handling editor was Renee Shellhaas, MD, MS.

Appendix Authors

Name	Location	Contribution
Amy Y. Tsou, MD, MSc	ECRI Evidence-based Practice Center, Plymouth Meeting, PA; Division of Neurology, Michael J Crescenzi Veterans Affairs Medical Center, Philadelphia	Drafting/revision of the manuscript for content, including medical writing for content; major role in the acquisition of data; study concept or design; and analysis or interpretation of data
Sudha Kilaru Kessler, MD, MSCE	Department of Pediatrics (Division of Neurology), Children’s Hospital of Philadelphia	Drafting/revision of the manuscript for content, including medical writing for content, and analysis or interpretation of data
Mingche Wu, MPH	ECRI Evidence based Practice Center, Plymouth Meeting, PA	Drafting/revision of the manuscript for content, including medical writing for content; major role in the acquisition of data; and analysis or interpretation of data
Nicholas S. Abend, MD, MSCE	Department of Pediatrics (Division of Neurology), Children’s Hospital of Philadelphia; Departments of Neurology and Pediatrics, University of Pennsylvania Perelman School of Medicine; Department of Anesthesia & Critical Care, University of Pennsylvania Perelman School of Medicine; Department of Biostatistics, Epidemiology and Informatics, University of Pennsylvania Perelman School of Medicine	Drafting/revision of the manuscript for content, including medical writing for content; study concept or design; and analysis or interpretation of data

Appendix (continued)

Name	Location	Contribution
Shavonne L. Massey, MD, MSCE	Department of Pediatrics (Division of Neurology), Children’s Hospital of Philadelphia; Departments of Neurology and Pediatrics, University of Pennsylvania Perelman School of Medicine	Drafting/revision of the manuscript for content, including medical writing for content; major role in the acquisition of data; and analysis or interpretation of data
Jonathan R. Treadwell, PhD	ECRI Evidence based Practice Center, Plymouth Meeting, PA;	Drafting/revision of the manuscript for content, including medical writing for content; major role in the acquisition of data; study concept or design; and analysis or interpretation of data

References

- Berg AT, Wusthoff C, Shellhaas RA, et al. Immediate outcomes in early life epilepsy: a contemporary account. *Epilepsy Behav.* 2019;97:44-50. doi: 10.1016/j.yebeh.2019.05.011.
- Wirrell E, Wong-Kissel L, Mandrekar J, Nickels K. Predictors and course of medically intractable epilepsy in young children presenting before 36 months of age: a retrospective, population-based study. *Epilepsia.* 2012;53(9):1563-1569.
- Shellhaas RA, Berg AT, Grinspan ZM, et al. Initial treatment for nonsyndromic early-life epilepsy: an unexpected consensus. *Pediatr Neurol.* 2017;75:73-79.
- Arts WFM, Brouwer OF, Peters AC, et al. Course and prognosis of childhood epilepsy: 5-year follow-up of the Dutch study of epilepsy in childhood. *Brain.* 2004;127(8):1774-1784.
- Beatty CW, Lockrow JP, Gedela S, Gehred A, Ostendorf AP. The missed value of underutilizing pediatric epilepsy surgery: a systematic review. *Semin Pediatr Neurol.* 2021;39:100917.
- Samanta D, Ostendorf AP, Willis E, et al. Underutilization of epilepsy surgery: part I: a scoping review of barriers. *Epilepsy Behav.* 2021;117:107837.
- Dwivedi R, Ramanujam B, Chandra PS, et al. Surgery for drug-resistant epilepsy in children. *N Engl J Med.* 2017;377(17):1639-1647.
- Buttle SG, Muir K, Dehnoei S, Webster R, Tu A. Referral practices for epilepsy surgery in pediatric patients: a North American Study. *Epilepsia.* 2022;63(1):86-95.
- Epilepsies: Diagnosis and Management.* National Institute for Health and Care Excellence. January 11, 2012. Updated May 2021. Accessed February 7, 2022. [guidelines.co.uk/neurology/nice-epilepsy-guideline/234875.article](https://www.nice.org.uk/neurology/nice-epilepsy-guideline/234875.article)
- Management of Infantile Epilepsies: Systematic Review.* Agency for Healthcare Research and Quality; October 25, 2012. [effectivehealthcare.ahrq.gov/products/management-infantile-epilepsy/research](https://www.effectivehealthcare.ahrq.gov/products/management-infantile-epilepsy/research).
- Methods Guide for Effectiveness and Comparative Effectiveness Reviews.* Effective Health Care Program, Agency for Healthcare Research and Quality; 2018. Last reviewed January 2020. Accessed February 23, 2022. [effectivehealthcare.ahrq.gov/products/collections/ceer-methods-guide](https://www.effectivehealthcare.ahrq.gov/products/collections/ceer-methods-guide)
- Wieser HG, Blume WT, Fish D, et al. Proposal for a new classification of outcome with respect to epileptic seizures following epilepsy surgery. *Epilepsia.* 2008;42(2):282-286.
- Engel J Jr, Van Ness PC, Rasmussen TB, Ojemann LM. *Outcome with Respect to Epileptic Seizures*; 1993:609-621.
- Sterne JAC, Savovic J, Page MJ, et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ.* 2019;366:4898.
- Sterne JA, Hernán MA, Reeves BC, et al. ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ.* 2016;355:i4919.
- Viswanathan M, Patnode C, Berkman ND, et al. *Assessing the Risk of Bias in Systematic Reviews of Health Care Interventions.* In: *Methods Guide for Comparative Effectiveness Reviews.* Agency for Healthcare Research and Quality; December 2017. Accessed February 7, 2022. [effectivehealthcare.ahrq.gov/products/methods](https://www.effectivehealthcare.ahrq.gov/products/methods).
- Berkman ND, Lohr KN, Ansari M, et al. *Grading the Strength of a Body of Evidence when Assessing Health Care Interventions for the Effective Health Care Program of the Agency for Healthcare Research and Quality: An Update.* Agency for Healthcare Research and Quality; 2013. Accessed February 7, 2022. [effectivehealthcare.ahrq.gov/topics/methods-guidance-grading-evidence/methods](https://www.effectivehealthcare.ahrq.gov/topics/methods-guidance-grading-evidence/methods).
- Reinholdson J, Olsson I, Edelvik A, et al. Long-term follow-up after epilepsy surgery in infancy and early childhood - a prospective population based observational study. *Seizure.* 2015;30:83-89.
- Kadish NE, Bast T, Reuner G, et al. Epilepsy surgery in the first 3 years of life: predictors of seizure freedom and cognitive development. *Neurosurgery.* 2019;84(6):E368-E377.
- Schramm J, Kuczaty S, Sassen R, Elger CE, Von Lehe M. Pediatric functional hemispherectomy: outcome in 92 patients. *Acta Neurochir (Wien).* 2012;154(11):2017-2028.

21. Kalbhenn T, Cloppenborg T, Wörmann FG, et al. Operative posterior disconnection in epilepsy surgery: experience with 29 patients. *Epilepsia*. 2019;60(9):1973-1983.
22. Iwasaki M, Uematsu M, Osawa SI, et al. Interhemispheric vertical hemispherotomy: a single center experience. *Pediatr Neurosurg*. 2015;50(5):295-300.
23. Otsuki T, Honda R, Takahashi A, et al. Surgical management of cortical dysplasia in infancy and early childhood. *Brain Develop*. 2013;35(8):802-809.
24. Sugimoto T, Otsubo H, Hwang PA, Hoffman HJ, Jay V, Iii OCS. Outcome of epilepsy surgery in the first three years of life. *Epilepsia*. 1999;40(5):560-565.
25. Steinbok P, Gan PYC, Connolly MB, et al. Epilepsy surgery in the first 3 years of life: a Canadian survey. *Epilepsia*. 2009;50(6):1442-1449.
26. Lettori D, Battaglia D, Sacco A, et al. Early hemispherectomy in catastrophic epilepsy. *Seizure* 2008;17(1):49-63.
27. Gaggero R, Consales A, Fazzini F, et al. Epilepsy associated with supratentorial brain tumors under 3 years of life. *Epilepsy Res*. 2009;87(2-3):184-189.
28. Roth J, Constantini S, Ekstein M, et al. Epilepsy surgery in infants up to 3 months of age: safety, feasibility, and outcomes: a multicenter, multinational study. *Epilepsia*. 2021;62(8):1897-1906.
29. Cook SW, Nguyen ST, Hu B, et al. Cerebral hemispherectomy in pediatric patients with epilepsy: comparison of three techniques by pathological substrate in 115 patients. *J Neurosurg Pediatr*. 2004;100(2):125-141.
30. Jonas R, Nguyen S, Hu B, et al. Cerebral hemispherectomy: hospital course, seizure, developmental, language, and motor outcomes. *Neurology*. 2004;62(10):1712-1721.
31. Kumar RM, Koh S, Knupp K, Handler MH, O'Neill BR. Surgery for infants with catastrophic epilepsy: an analysis of complications and efficacy. *Childs Nerv Syst*. 2015;31(9):1479-1491.
32. Loddenkemper T, Holland KD, Stanford LD, Kotagal P, Bingaman W, Wyllie E. Developmental outcome after epilepsy surgery in infancy. *Pediatrics* 2007;119(5):930-935. doi: 10.1542/peds.2006-2530
33. Pinto ALR, Lohani S, Bergin AMR, et al. Surgery for intractable epilepsy due to unilateral brain disease: a retrospective study comparing hemispherectomy techniques. *Pediatr Neurol*. 2014;51(3):336-343. doi: 10.1016/j.pediatrneurol.2014.05.018
34. Maton B, Jayakar P, Resnick T, Morrison G, Ragheb J, Duchowny M. Surgery for medically intractable temporal lobe epilepsy during early life. *Epilepsia*. 2008;49(1):80-87. doi: 10.1111/j.1528-1167.2007.01315.x
35. Dunkley C, Kung J, Scott RC, et al. Epilepsy surgery in children under 3 years. *Epilepsy Res*. 2011;93(2-3):96-106. doi: 10.1016/j.eplesyres.2010.11.002
36. Iwasaki M, Iijima K, Kawashima T, et al. Epilepsy surgery in children under 3 years of age: surgical and developmental outcomes. *J Neurosurg Pediatr*. 2021;28(4):395-403. doi: 10.3171/2021.4.peds21123
37. Perry MS, Shandley S, Perelman M, et al. Surgical evaluation in children <3 years of age with drug-resistant epilepsy: patient characteristics, diagnostic utilization, and potential for treatment delays. *Epilepsia*. 2022;63(1):96-107.
38. Jobst BC, Cascino GD. Resective epilepsy surgery for drug-resistant focal epilepsy: a review. *JAMA*. 2015;313(3):285-293. doi: 10.1001/jama.2014.17426
39. England MJ, Liverman CT, Schultz AM, Strawbridge LM. Epilepsy across the spectrum: promoting health and understanding. *Epilepsy Behav*. 2012;25(2):266-276. doi: 10.1016/j.yebeh.2012.06.016
40. Donahue MA, Herman ST, Dass D, et al. Establishing a learning healthcare system to improve health outcomes for people with epilepsy. *Epilepsy Behav*. 2021;117:107805.

Neurology®

Surgical Treatments for Epilepsies in Children Aged 1–36 Months: A Systematic Review

Amy Y. Tsou, Sudha Kilaru Kessler, Mingche Wu, et al.
Neurology 2023;100:e1-e15 Published Online before print October 21, 2022
DOI 10.1212/WNL.0000000000201012

This information is current as of October 21, 2022

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/100/1/e1.full
References	This article cites 34 articles, 3 of which you can access for free at: http://n.neurology.org/content/100/1/e1.full#ref-list-1
Citations	This article has been cited by 1 HighWire-hosted articles: http://n.neurology.org/content/100/1/e1.full##otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Epilepsy surgery http://n.neurology.org/cgi/collection/epilepsy_surgery_
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2022 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

