

SARS-CoV-2 Vaccination and Autoimmune Neuropathies

Rebalancing the Risk

Jeffrey A. Allen, MD

Neurology® 2023;100:55-56. doi:10.1212/WNL.000000000201366

With every influenza vaccination, the question is asked: “Have you ever had Guillain-Barré syndrome?” If so, the potential vaccination recipient is advised to talk to their doctor before proceeding, and the physician will invariably recommend against it. The topic is widely discussed in patient and physician networks, with stakeholders holding firm convictions for or (more commonly) against vaccination. Naturally, with the emergence of the COVID-19 pandemic and the reality of administering billions of SARS-CoV-2 vaccinations across the globe, the anxiety within the inflammatory neuropathy community and among clinicians caring for patients with GBS and CIDP has been raised to new heights.

But why? What fuels this anxiety? In 1976, a looming swine influenza pandemic prompted the development of a national vaccination campaign program within the United States.¹ In its wake, an 8-fold increase in postvaccination GBS cases was observed (about 1 GBS case per 100,000 vaccinations), and the Institute of Medicine concluded that a causal relationship between the 1976 swine influenza vaccine and GBS was likely.² Coupled with the fact that the swine flu pandemic of 1976 never materialized, the foundation for vaccination hesitancy among patients with GBS/CIDP and their physicians was formed. However, since 1976, associations between seasonal vaccinations and GBS have been meager at best. Most studies have shown no increased risk³ while others showed a marginal increased risk on the order of 1–2 additional cases of GBS per 1 million persons vaccinated.⁴ To give these numbers more context, consider the stronger association observed between influenza infections and GBS where an estimated risk of about 17 GBS cases per 1 million patients hospitalized with influenza have been reported.⁵

Now enter the COVID-19 pandemic. Although much is still to be learned, there appears to be a small increased risk of GBS after COVID-19 infection⁶ and probably an even smaller increase risk after some (but not all) of the SARS-CoV-2 vaccinations. GBS after the adenovirus vaccinations ChAdOx1 nCoV-19 (AstraZeneca) and Ad26.COV2 (Johnson & Johnson) ranges between 4–7 cases per million vaccinations.^{7,8} No increased risk has been observed after the mRNA tozinameran (Pfizer-BioNTech) or 1 mRNA-1273 (Moderna) vaccines.⁸

So what is the patient with GBS/CIDP/MMN to do when faced with vaccination in the time of COVID-19? The threat of voluntarily doing something to worsen their disease can be terrifying, but so too is the prospect of developing a severe infection. Extrapolating from the influenza vaccine experience, patients are primed to be wary of vaccines. The information conveyed by support networks and sometimes physicians is often not reassuring and frequently inaccurate because it fails to balance risks of triggering relapse from vaccination with the risks from actually becoming infected.

In this issue of *Neurology*®, Baars et al. take on the timely issue of SARS-CoV-2 vaccination safety in patients with preexisting inflammatory neuropathies.⁹ This prospective, multicenter cohort study included 406 patients with confirmed GBS, MMN, or CIDP who received a SARS-CoV-2 vaccination. Although disability and quality-of-life scales were collected, deterioration was loosely defined as any self-reported change in disease-related symptoms. Assessments were conducted 48 hours before vaccination and within 6 weeks to 4 months after vaccination.

Correspondence

Dr. Allen
jaallen@umn.edu

RELATED ARTICLE

Research Article

SARS-CoV-2 Vaccination Safety in Guillain-Barré Syndrome, Chronic Inflammatory Demyelinating Polyneuropathy, and Multifocal Motor Neuropathy

Page 74

MORE ONLINE

COVID-19 Resources

For the latest articles, invited commentaries, and blogs from physicians around the world

[NPub.org/COVID19](https://www.npub.org/COVID19)

From the University of Minnesota, Minneapolis, MN.

Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the editorial.

Symptoms experienced within 48 hours after vaccination were classified as vaccination side effects (i.e., not related to the immune-mediated neuropathy).

Among 162 patients with GBS, none experienced a GBS recurrence. Although 12 of 241 patients with CIDP or MMN reported worsening disease-related symptoms, the change prompted treatment modifications in only 6 patients, and in all, the symptoms resolved within 8 weeks. Also notably, in 10 of these 12 patients, other explanations for symptom exacerbation were present during or leading into the vaccination.

These results should provide much needed reassurance to inflammatory neuropathy patients in need of SARS-CoV-2 vaccination. In the unusual event that symptoms worsened, in most instances the setback was mild and self-limiting. Digging a bit further, a notable study limitation was how deterioration was defined. Objective documentation of neuropathy worsening with measures such as grip strength testing or neurologic examination was not feasible, and instead, there was a reliance on self-reported symptom changes to capture deterioration. Generally speaking, placebo and nocebo responses driven by anticipation and expectation are common in CIDP during changes in treatment or in the context of other life events.¹⁰ This phenomena likely explains some of the self-reported deterioration in patients with CIDP and MMN observed in this study, rather than worsening of the neuropathy. In only 3 patients, did the change in disability scores reach the minimal clinically important difference, and in 10 of 12 patients, other explanations were observed that may explain the change in symptom status. Just as objectification of inflammatory neuropathy relapse was a challenge in this study, so too is it a struggle during clinical practice. Although it may be attractive to attribute any change in the postvaccination period to worsening neuropathy, it is critical to objectify changes and identify confounding explanations before assigning causation.

Extremely rare events are challenging to capture in relatively small prospective studies of this kind. It is even more difficult to understand if the mRNA and adenovirus vaccinations confer different levels of risk, as seems to be the case in the general population.^{7,8} Although this study is unable to exclude

associations between SARS-CoV-2 vaccination and worsening inflammatory neuropathy, it can now be confidently state that if there is a risk it is likely to be very small and almost certainly less than the risk of an infection-triggered relapse. The time is now to rebalance the risk of vaccination in our patients with GBS, CIDP, and MMN. It is overdue for influenza vaccination. Let us not make the same mistake with COVID-19.

Study Funding

The author reports no targeted funding.

Disclosure

J.A. Allen provides consultant services to Argenx, Alexion, Annexon, Axcea, CSL Behring, Takeda, Grifols, Pfizer, Johnson & Johnson. Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures.

Publication History

Received by *Neurology* July 31, 2022. Accepted in final form August 16, 2022.

References

1. Schonberger LB, Bregman DJ, Sullivan-Bolyai JZ, et al. Guillain-Barre syndrome following vaccination in the national influenza immunization program, United States, 1976-1977. *Am J Epidemiol.* 1979;110(2):105-123.
2. Langmuir AD, Bregman DJ, Kurland LT, Nathanson N, Victor M. An epidemiologic and clinical-evaluation of Guillain-Barre-syndrome reported in association with the administration of swine influenza vaccines. *Am J Epidemiol.* 1984;119(6): 841-879.
3. Baxter R, Bakshi N, Fireman B, Lewis B, Ray P, et al. Lack of association of Guillain-Barre syndrome with vaccinations. *Clin Infect Dis.* 2013;57(2):197-204.
4. Juurlink DN, Stukel TA, Kwong J, Kopp McGreerA, Upshur R, et al. Guillain-Barre syndrome after influenza vaccination in adults—a population-based study. *Arch Intern Med.* 2006;166(20):2217-2221.
5. Sellers SA, Hagan RS, Hayden FG, Fischer WA II. The hidden burden of influenza: a review of the extra-pulmonary complications of influenza infection. *Influenza Other Respir Viruses.* 2017;11:372-393.
6. Luijten L, Leonhard S, van der Eijk A, et al. Guillain-Barré syndrome after SARS-CoV-2 infection in an international prospective cohort study. *Brain* 2021;144(11): 3392-3404.
7. Keh R, Scanlon SM, Datta-Nemdhary P, Donegan K, Cavangh S, Foster M, et al. COVID-19 vaccination and Guillain-Barré syndrome: analyses using the National Immunoglobulin Database. *Brain.* 2022;awac067. doi: 10.1093/brain/awac067. Online ahead of print.
8. Woo EJ, Mba-Jones A, Dimova R, Alimchandani M, Zinderman C, Nair N., Association of receipt of the Ad26.COV2.S COVID-19 vaccine with presumptive Guillain-Barré syndrome, february-july 2021. *JAMA* 2021;326(16):1606-1613.
9. Baaes AE, Kuitwaard K, de Koning LC, et al. Safety of SARS-CoV-2 vaccination in Guillain-Barré syndrome, chronic inflammatory demyelinating polyneuropathy and multifocal motor neuropathy. *Neurology* 2023;100:e182-e191.
10. Lewis RA, Cornblath DR, Hartung HP, et al. Placebo effect in chronic inflammatory demyelinating polyneuropathy: the PATH study and a systematic review. *J Peripher Nerv Syst.* 2020;25(3):230-237.

Neurology[®]

SARS-CoV-2 Vaccination and Autoimmune Neuropathies: Rebalancing the Risk

Jeffrey A. Allen

Neurology 2023;100;55-56 Published Online before print September 20, 2022

DOI 10.1212/WNL.0000000000201366

This information is current as of September 20, 2022

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/100/2/55.full
References	This article cites 10 articles, 1 of which you can access for free at: http://n.neurology.org/content/100/2/55.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Immunology http://n.neurology.org/cgi/collection/all_immunology Chronic inflammatory demyelinating polyneuropathy http://n.neurology.org/cgi/collection/chronic_inflammatory_demyelinating_polyneuropathy Guillain-Barre syndrome http://n.neurology.org/cgi/collection/guillainbarre_syndrome Peripheral neuropathy http://n.neurology.org/cgi/collection/peripheral_neuropathy
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2022 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

