

Teaching NeuroImage: Bloomy Rind Sign of Leptomeningeal Carcinomatosis


Xi Liu, PhD, MD,* Changhong Tan, PhD, MD,* Yuxia Zhu, MD, Sichen Li, MD, Hongyu Lai, Fen Deng, MD, and Lifen Chen, PhD, MD

Neurology® 2023;101:e99-e100. doi:10.1212/WNL.0000000000207124

Correspondence

Dr. Chen
lifen_chen@163.com

Figure 1 Symmetrical Curvilinear Abnormality Along Surface of Midbrain (A), Pons (B), and Cerebellar Peduncles (C) in Axial Brain MRI Images, FLAIR Sequences at Admission (Arrows), While Medulla Was Relatively Intact (D)



A 68-year-old woman presented with vertigo, nausea, and vomiting for 3 days. She had lung adenocarcinoma with *EGFR* mutation (p.L858R) for 1 year and was in remission after osimertinib treatment. Neurologic examination revealed stiff neck, horizontal and vertical nystagmus, dysphagia, and bilateral Babinski signs. MRI revealed a symmetrical hyperintensity along the surface of the midbrain, pons, and cerebellar peduncles in T2, fluid attenuated inversion recovery, and diffusion-weighted imaging (Figure 1). T1 and apparent diffusion coefficient were normal, and there was no contrast enhancement. No significant supratentorial anomaly was found. CSF cytology identified cancer cells carrying *EGFR* mutation (p.L858R). We added bevacizumab to osimertinib. MRI 2 months later revealed thickened lesions expanded to the medulla and inner ears with hydrocephalus (Figure 2).

MORE ONLINE

Teaching slides


links.lww.com/WNL/C649

*Xi Liu and Changhong Tan contributed equally to this work as first authors.

From the Department of Neurology, The Second Affiliated Hospital of Chongqing Medical University, China.

Go to Neurology.org/N for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article.

Figure 2 Thickened Lesions Along Surface of Midbrain (A), Pons (B), and Cerebellar Peduncles (C) (Arrows), Which Expanded to Medulla and Inner Ears (D) (Asterisks) in Axial Brain MRI Images, Fluid Attenuated Inversion Recovery Sequences at Follow-up



The patient finally died because of respiratory failure. In leptomeningeal carcinomatosis, common MRI findings are linear enhancement of the sulci, leptomeninges, and nerve roots, whereas bloomy rind sign, possibly induced by tumor infiltration, cytotoxic edema, and microinfarction, is rare and characteristic.^{1,2}

Author Contributions

X. Liu: drafting/revision of the manuscript for content, including medical writing for content; major role in the acquisition of data; study concept or design; analysis or interpretation of data. C. Tan: drafting/revision of the manuscript for content, including medical writing for content; major role in the acquisition of data. Y. Zhu: major role in the acquisition of data. S. Li: major role in the acquisition of data. H. Lai: drafting/revision of the manuscript for content, including medical writing for content. F. Deng: major role in the acquisition of data; analysis or interpretation of data. L. Chen: major role in the acquisition of data; study concept or design; analysis or interpretation of data.

Study Funding

National Natural Science Foundation of China (82001367).

Disclosure

X. Liu reports that he received funding from National Natural Science Foundation of China (82001367) for supporting this manuscript. The other authors report no relevant disclosures. Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures.

Publication History

Received by *Neurology* October 3, 2022. Accepted in final form January 12, 2023. Submitted and externally peer reviewed. The handling editor was Resident and Fellow Section Editor Whitley Aamodt, MD, MPH.

References

1. Crombe A, Alberti N, Durieux M, Frulio N, Dousset V, Tourdias T. Exceptional symmetric anterior brainstem involvement in leptomeningeal carcinomatosis. *J Neuroradiol*. 2014;41(4):279-281.
2. Mitsuya K, Nakasu Y, Deguchi S, et al. FLAIR hyperintensity along the brainstem surface in leptomeningeal metastases: a case series and literature review. *Cancer Imaging*. 2020;20(1):84.

Neurology®

Teaching NeuroImage: Bloomy Rind Sign of Leptomeningeal Carcinomatosis

Xi Liu, Changhong Tan, Yuxia Zhu, et al.

Neurology 2023;101:e99-e100 Published Online before print February 16, 2023

DOI 10.1212/WNL.0000000000207124

This information is current as of February 16, 2023

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/101/1/e99.full
References	This article cites 2 articles, 0 of which you can access for free at: http://n.neurology.org/content/101/1/e99.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Metastatic tumor http://n.neurology.org/cgi/collection/metastatic_tumor MRI http://n.neurology.org/cgi/collection/mri
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2023 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

